

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 0 650 965 B1

(12) EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
07.02.2001 Bulletin 2001/06

(51) Int Cl.7: C07D 333/20, C07F 9/06,  
C07C 53/10, C07C 63/08

(21) Application number: 94307394.0

(22) Date of filing: 07.10.1994

(54) **Asymmetric synthesis of  
(S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine an intermediate in the  
preparation of duloxetine**

Asymmetrische Synthese von (S)-(+)-N,N-Dimethyl-3-(1-Naphthyloxy)-3-(2-Thienyl)propanamin  
einem Zwischenprodukt in der Synthese von Duloxetin

Synthèse asymétrique de (S)-(+)-N,N-diméthyl-3-(1-naphtalényloxy)-3-(2-thiényl)propénamine un  
produit intermédiaire pour la préparation de duloxetine

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IE IT LI LU NL PT  
SE  
Designated Extension States:  
LT SI

(30) Priority: 12.10.1993 US 135032

(43) Date of publication of application:  
03.05.1995 Bulletin 1995/18

(73) Proprietor: ELI LILLY AND COMPANY  
Indianapolis, Indiana 46285 (US)

(72) Inventor: Berglund, Richard Alan  
Lafayette, Indiana 47905 (US)

(74) Representative: Hudson, Christopher Mark et al  
Lilly Industries Limited  
European Patent Operations  
Erl Wood Manor  
Windlesham Surrey GU20 6PH (GB)

(56) References cited:  
EP-A- 0 273 658 EP-A- 0 391 070  
EP-A- 0 457 559 EP-A- 0 509 241  
WO-A-94/00416 DE-A- 4 123 253  
US-A- 4 558 164

- TETRAHEDRON LETTERS, vol.31, no.49, 26 November 1990 pages 7101 - 7104 DEETER J. ET AL. 'Asymmetric synthesis and absolute stereochemistry of LY248686'
- TETRAHEDRON: ASYMMETRY, vol.3, no.4, 1992 pages 525 - 528 SCHNEIDER M.P. & GOERGENS U. 'An efficient route to enantiomerically pure antidepressants: Tomoxetine, nisoxetine and fluoxetine'
- CHEMICAL ABSTRACTS, vol. 116, no. 21, 25 May 1992, Columbus, Ohio, US; abstract no. 214138p, YOSHIDA N. & MIYAZAWA K. 'Preparation of optically active fluoxetine and N-methyl-3-phenyl-3-hydroxypropionamide' page 706 ; column 1 ; & JP-A-04 005 268 (CHISSO CORP.) 9 January 1992
- JOURNAL OF ORGANIC CHEMISTRY, vol.53, no.17, 1988 pages 4081 - 4084 GAO Y. & SHARPLESS K.B. 'Asymmetric synthesis of both enantiomers of tomoxetine and fluoxetine, Selective reduction of 2,3-epoxycinnamyl alcohol with Red-Al'

Remarks:


The file contains technical information submitted after the application was filed and not included in this specification

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

[0001] This invention belongs to the fields of pharmaceutical chemistry and synthetic organic chemistry, and provides an asymmetric process for the synthesis of a key intermediate in the preparation of duloxetine, (+)-N-methyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine, hydrochloric acid salt.

[0002] Duloxetine is a pharmaceutical now under development as an anti-depressant. It inhibits the uptake of both norepinephrine and serotonin and is presently in clinical evaluation. The compound was disclosed in U.S. Patent Nos. 5,023,269 and 4,956,388 by Robertson, *et al.*, and the synthesis of it was discussed in more detail by Deeter, *et al.*, in *Tetrahedron Letters*, 31(49), 7101-04 (1990). The process for preparing duloxetine is there outlined as follows.



[0003] The product of Step D above, when in the hydrochloride salt form, is duloxetine.

[0004] The present invention provides improved conditions for carrying out Step C in the above scheme, whereby the product of Step C is obtained more quickly in better purity and yield than has previously been possible.

[0005] A particularly useful salt for the isolation of the intermediate is also provided.

[0006] The present invention provides a process for preparing (S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine comprising reacting (S)-(-)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropanamine with sodium hydride, potassium benzoate or potassium acetate, and 1-fluoronaphthalene in an organic solvent; and if desired, recovering the product as the phosphoric acid salt. The phosphoric acid salt is also an aspect and an embodiment of the invention.

[0007] The present invention provides an asymmetric process for preparing the specific enantiomer shown above as the product of Step C in the scheme. It is named as the (S)-(+) enantiomer and that nomenclature will be used throughout this document. The starting material for the process of this invention, the hydroxypropanamine, is named as the (S)-(-) enantiomer.

[0008] The preferred solvent for carrying out the present invention is dimethylsulfoxide. Other solvents, such as dimethylformamide or pyridine, may be used as well. The process may be effectively carried out at unusually high concentrations, such as from 0.5-1.0 molar, preferably 0.6-0.9 molar. However, the rate and yield of the process is not seriously affected by the concentration, so long as the solubility limit is not exceeded, of course.

[0009] The order and manner of combining the reactants are not important and may be varied. The reactants may be added to the reaction mixture as solids, or may be dissolved individually and combined as solutions. Further any of the reactants may be dissolved together as subgroups, and those solutions may be combined in any order.

[0010] A preferred manner of carrying out the invention is to dissolve the starting material and sodium hydride together, add the potassium compound, and then add the 1-fluoronaphthalene, and that embodiment will be further described in detail.

[0011] When the starting material has been dissolved, preferably at ambient temperature such as from 10° to 35°C,

a portion of sodium hydride is added. The amount of sodium hydride is equimolar with the starting material; no particular advantage is found in using excess sodium hydride. The reaction mixture is then stirred for a period, such as from 5 to 60 minutes, and then combined with potassium benzoate or potassium acetate and 1-fluoronaphthalene.

[0012] Only a small amount of the potassium compound is needed, from 0.05 to 1 equivalent. In general, the advantages of the present invention are best obtained when only from 0.1 to 0.3 equivalent of the potassium compound is used. There is no disadvantage in using more, except the obvious cost disadvantage.

[0013] An additional period of stirring is provided after the addition of the potassium compound, and then the 1-fluoronaphthalene is added. A small amount of excess 1-fluoronaphthalene, such as from 1% to 25% excess, may advantageously be used to assure consumption of the more expensive thiophene starting material.

[0014] When the fluoronaphthalene has been added, the reaction mixture is preferably warmed, to a temperature from 40° to 75°C, preferably from 45° to 70°C, and most preferably from 60° to 65°C, and the mixture is stirred for a short period of time, such as from 1 to 5 hours, most preferably from 1.5 to 4 hours.

[0015] The desired product is then isolated by conventional extractions and filtrations, and, if it is desired to do so, the product may advantageously be converted to the phosphoric acid salt by reaction with phosphoric acid in an organic solvent such as ethyl acetate.

[0016] The advantage of the present invention is found in its ability to prepare the desired product in yields in the range of 95%, with very little racemization, in short periods of time as described above. Previous procedures took more than a day, and gave product of inferior purity.

[0017] The hydroxypropanamine which is the starting material for the present process is available by the teachings of the prior art, but preparation 1 below is provided to assure its availability to the reader of the present document.

[0018] The product of the present process, the dimethyl compound, is converted to duloxetine by demethylating to provide the desired monomethyl pharmaceutical, and conversion to the hydrochloride salt. Preparation 2 below illustrates that conversion, which, like the preparation of the hydroxy starting material, is from the prior art and forms no part of the present invention.

#### **PREPARATION 1**

(S)-(-)-N,N-dimethyl-3-hydroxy-3-(2-thienyl)-propanamine

[0019] A mixture of 8.18 g of 2-acetylthiophene, 6.66 g of dimethylamine hydrochloride, 2.9 g of paraformaldehyde and 0.31 g of concentrated hydrochloric acid in 20 ml of isopropanol was heated to reflux and stirred for 6 hours. The mixture was then cooled to 0°C and stirred for one hour more. The slurry was then filtered, and the solid was washed with cold ethanol. The washed solid was dried for 16 hours at 50°C to obtain 12.5 g of 2-thienyl 2-dimethylaminoethyl ketone hydrochloride, as a white solid. A 12.0 g portion of that intermediate product was stirred in 40 ml of ethanol at ambient temperature, and the pH of the solution was raised to 11-12 by slow addition of sodium hydroxide. A 1.03 g portion of sodium borohydride was added, and the mixture was stirred at ambient temperature for 4 hours. Then 7.5 ml of acetone was added, and the mixture was stirred for 20 minutes more. The mixture was then concentrated by evaporation to a white slurry, and 120 ml of methyl t-butyl ether was added. The mixture was acidified to pH 1-1.5 by addition of concentrated hydrochloric acid, and the solution was stirred for ten minutes. The pH was then made basic to pH 12 by slow addition of sodium hydroxide.

[0020] The layers were then separated, the aqueous phase was extracted with 30 ml of methyl t-butyl ether, and the organic phases were combined and washed once with 50 ml of water. The organic phase was concentrated by evaporation to 118 ml, and was heated to 50°C.

[0021] In a separate vessel, (S)-(+)-mandelic acid, 4.18 g, was dissolved in 12 ml of ethanol at 50°C, and the mandelic acid solution was added slowly to the previous solution. The resulting slurry was then heated to reflux and stirred for 45 minutes. It was then cooled to ambient temperature, stirred for one hour, and filtered, and the solid was washed with methyl t-butyl ether. The solid was then dried under vacuum at 50°C to obtain 7.29 g of the mandelic acid salt of the desired product, which is isolated as the free amine by dissolution in water, basification with sodium hydroxide solution, extraction into an organic solvent, and evaporation to remove the solvent.

#### **EXAMPLE 1**

(S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine, phosphoric acid salt

[0022] A 13.5 g portion of (S)-(-)-N,N-dimethyl-3-hydroxy-3-(2-thienyl)propanamine was dissolved in 80 ml of dimethylsulfoxide at 25°C. To the solution was slowly added 3 g of sodium hydride as a 60% dispersion in mineral oil, with vigorous stirring. After 15 minutes of stirring, 1.17 g of potassium benzoate was added and stirring was continued at approximately constant temperature for another 15 minutes. Then, 12.8 g of 1-fluoronaphthalene was slowly added to

the reaction mixture, and after the addition was complete, the mixture was heated and was stirred for 2.5 hours at 60-65°C. The mixture was then poured slowly into 190 ml of cold water and the pH was adjusted to 4.8 by addition of acetic acid. The temperature of the mixture was brought to 25°C, and 75 ml of hexane was added and stirring was continued for 10 minutes. The layers were then separated and the aqueous phase was stirred again with 75 ml of hexane and the phases separated. The pH of the aqueous phase was adjusted to 10.2 by addition of aqueous sodium hydroxide, and 75 ml of ethyl acetate was added. That mixture was stirred for 15 minutes at 25°C, and the 2-phase mixture was vacuum filtered through a pad of filter aid. The phases of the filtrate were allowed to separate, and the aqueous phase was extracted with 75 ml of ethyl acetate. The extract was combined with the previous ethyl acetate layer, and that mixture was washed with 100 ml of water. The organic layer was stirred at 25°C, and to it was added, dropwise, 7 g of 85% phosphoric acid. After the addition was complete, the mixture was stirred for 20 minutes more and was then cooled to 0°C and stirred for 1 hour at that temperature. The slurry was then filtered and the solids washed three times with 20 ml portions of cold ethyl acetate. The solid was dried at 60°C to afford 24.19 g of the title compound as a white solid, 98.1% potency, adjusted yield 79.6%, 91% EE.

## Assay Methodology

**[0023]** The product of the example was analyzed by high performance liquid chromatography, using a Spectra Physics SP 8800 instrument equipped with a SP 4400 integrator and a Spectroflow 757 detector, set at 230 nm, at a sensitivity of 0.5 absorption units, 1 second filter rise time. The column was a Dupont Zorbax Rx C8, 4.6 mm x 25 cm. The eluant was 70% acetonitrile, 30% 0.01 M phosphate buffer at pH 6, flow rate of 1.0 ml/minute, injection volume 20 microliters. The samples were prepared by diluting 0.1 to 0.3 g of reaction mixture or extract to 50 ml with 1:1 acetonitrile:water. The product peak elutes at 13-17 minutes; starting material at 6-8 minutes; fluoronaphthalene at 5-6 minutes; dimethylsulfoxide at 2-3 minutes; and potassium benzoate at 2-2.5 minutes.

**[0024]** When a chiral assay was to be done, the same equipment was set at 280 nm and a sensitivity of 0.1 absorption unit, and a Chiralcel OD column was used. The eluant for chiral assays was 2% isopropanol, 0.2% diethylamine, and 97.8% hexane. The same injection and flow settings were used. The samples were prepared by diluting 0.1-0.3 g of reaction mixture or extract to 5 ml with dichloromethane, washing the mixture with about 5 ml of water, and drying the organic phase over sodium sulfate. The resulting solution was filtered and diluted to 25 ml with eluant. The desired enantiomer elutes at 5-5.5 minutes, the undesired enantiomer at 6-6.5 minutes and fluoronaphthalene at 3-4 minutes.

## Example 2

(S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine

**[0025]** A 1.60 g portion of (S)-(+)-N,N-dimethyl-3-hydroxy-3-(2-thienyl)propanamine was dissolved in 8 ml of dimethylsulfoxide at ambient temperature, to which was added 0.35 g of sodium hydride as a 60% dispersion in mineral oil with vigorous stirring. After 30 minutes of stirring, 0.28 g of potassium benzoate was added, and stirring was continued for 10 minutes more. Then 1.52 g of 1-fluoronaphthalene was added and the mixture was then stirred at 50°C for 8 hours. The reaction mixture was poured slowly into 30 ml of cold water, and the pH was adjusted to 4.8 by addition of acetic acid. Fifteen ml of hexane was added, the mixture was stirred for 10 minutes, and the layers were separated. The aqueous phase was stirred again with 15 ml of hexane and the phases separated. The pH of the aqueous phase was adjusted to 12.5 by addition of aqueous sodium hydroxide, and 15 ml of ethyl acetate was added. The basic mixture was stirred at ambient temperature for 10 minutes, and the layers were separated. The aqueous phase was extracted with another 15 ml portion of ethyl acetate, and the organic extracts were combined, washed once with 30 ml of water, and dried over magnesium sulfate. The solvent was removed under vacuum to obtain a red-brown oil which was dissolved in the minimal amount of 1:1 ethyl acetate:hexane. The solution was flushed through a pad of silica gel using as eluant, ethyl acetate:hexane:methanol:ammonium hydroxide, 47:47:5.8:0.2. The product fraction was evaporated under vacuum to obtain 2.3 g of the desired product as an amber oil.

## Preparation 2

(S)-(+)-N-methyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine hydrochloride

**[0026]** Five g of the product of Example 1 was stirred in a mixture of 40 ml of toluene and 40 ml of water at 40°C, and 2.5 ml of 30% ammonium hydroxide solution was added. The mixture was stirred for 10 minutes at constant temperature and the layers were separated. The organic phase was washed with water, dried with magnesium sulfate and filtered. The filtrate was concentrated to half volume under vacuum and was heated to 55°C. Then 0.16 g of diisopropylethylamine was added, followed by the dropwise addition of 2.39 g of phenyl chloroformate. The mixture

was stirred at 55°C for 1.25 hours, and 50 ml of 1% sodium bicarbonate solution was added. The mixture was stirred for ten minutes at 40-50°C, and the phases were separated. The organic phase was washed twice with 0.5 N hydrochloric acid, and then washed with 1% sodium bicarbonate solution. The washed organic phase was divided in half, and one aliquot was evaporated under vacuum and 26 ml of dimethylsulfoxide was added to the residue. The mixture was heated to 45°C, and 1 g of sodium hydroxide and 6 ml of water was added dropwise. The basic mixture was stirred for 18 hours at 50°C, diluted with 17 ml of water, and acidified to pH 5.0-5.5 by addition of acetic acid. Then 20 ml of hexane was added, the mixture was stirred for ten minutes, and the phases separated. The aqueous phase was made basic to pH 10.5 by addition of 50% aqueous sodium hydroxide, and 17 ml of ethyl acetate was added. After stirring for 10 minutes, the phases were separated, and the aqueous layer was extracted with another 17 ml of ethyl acetate. The combined organic extracts were washed with water and concentrated to 10 ml under vacuum. 0.46 g of concentrated hydrochloric acid was added to the residue, and then a seed crystal and an additional 10 ml of ethyl acetate was added. The mixture was stirred for 30 minutes more, and the solution was concentrated to 10 ml under vacuum. The residue was stirred for 1 hour at ambient temperature and 1 hour at 0°C to produce a slurry, which was filtered. The solid was washed with chilled ethyl acetate to obtain 1.32 g of the desired product, which was duloxetine as a white solid of potency 99.8%.

[0027] The above Example illustrates the convenience and the excellent results of the present process.

## Claims

1. A process for preparing (S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine comprising reacting (S)-(-)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropanamine with sodium hydride, a potassium compound chosen from potassium benzoate or potassium acetate, and 1-fluoronaphthalene in an organic solvent.
2. A process of Claim 1 wherein the product is recovered as the phosphoric acid salt.
3. A process of Claim 1 wherein the (S)-(-)-N,N-dimethyl-3-(2-thienyl)-3-hydroxypropanamine is dissolved with the sodium hydride in the organic solvent, the potassium compound is then added with stirring and then the 1-fluoronaphthalene is added with additional stirring.
4. A process of Claim 3 wherein the product is isolated as the phosphoric acid salt.
5. A process of Claim 1 wherein the potassium compound is potassium benzoate.
6. A process of Claim 3 wherein the potassium compound is potassium benzoate.
7. A process of Claim 4 wherein the potassium compound is potassium benzoate.
8. (S)-(+)-N,N-dimethyl-3-(1-naphthalenyloxy)-3-(2-thienyl)propanamine, phosphoric acid salt.

## Patentansprüche

1. Verfahren zur Herstellung von (S)-(+)-N,N-Dimethyl-3-(1-naphthalinyloxy)-3-(2-thienyl)propanamin durch Umsetzen von (S)-(-)-N,N-Dimethyl-3-(2-thienyl)-3-hydroxypropanamin mit Natriumhydrid, einer Kaliumverbindung, die aus Kaliumbenzoat oder Kaliumacetat ausgewählt ist, und 1-Fluornaphthalin in einem organischen Lösungsmittel.
2. Verfahren nach Anspruch 1, wobei das Produkt in Form eines Phosphorsäuresalzes gewonnen wird.
3. Verfahren nach Anspruch 1, wobei das (S)-(-)-N,N-Dimethyl-3-(2-thienyl)-3-hydroxypropanamin mit dem Natriumhydrid in einem organischen Lösungsmittel gelöst wird, worauf die Kaliumverbindung unter Rühren zugegeben wird und anschließend das 1-Fluornaphthalin unter weiterem Rühren zugegeben wird.
4. Verfahren nach Anspruch 3, wobei das Produkt in Form des Phosphorsäuresalzes isoliert wird.
5. Verfahren nach Anspruch 1, wobei die Kaliumverbindung Kaliumbenzoat ist.
6. Verfahren nach Anspruch 3, wobei die Kaliumverbindung Kaliumbenzoat ist.

## EP 0 650 965 B1

7. Verfahren nach Anspruch 4, wobei die Kaliumverbindung Kaliumbenzoat ist.
8. (S)-(+)-N,N-Dimethyl-3-(1-naphthalinyloxy)-3-(2-thienyl)propanaminphosphorsäuresalz.

5

### Revendications

1. Procédé de préparation de (S)-(+)-N,N-diméthyl-3-(1-naphtalényloxy)-3-(2-thiényl)propanamine comprenant la réaction de (S)-(-)-N,N-diméthyl-3-(2-thiényl)-3-hydroxypropanamine avec de l'hydrure de sodium, un composé de potassium, choisi parmi le benzoate de potassium ou l'acétate de potassium et du 1-fluoronaphtalène dans un solvant organique.
2. Procédé selon la revendication 1, dans lequel le produit est récupéré sous forme du sel d'acide phosphorique.
3. Procédé selon la revendication 1, dans lequel la (S)-(-)-N,N-diméthyl-3-(2-thiényl)-3-hydroxypropanamine est dissoute avec l'hydrure de sodium dans le solvant organique, le composé de potassium y est ensuite ajouté sous agitation, puis le 1-fluoronaphtalène y est ajouté sous agitation supplémentaire.
4. Procédé selon la revendication 3, dans lequel le produit est isolé sous forme du sel d'acide phosphorique.
5. Procédé selon la revendication 1, dans lequel le composé de potassium est le benzoate de potassium.
6. Procédé selon la revendication 3, dans lequel le composé de potassium est le benzoate de potassium.
7. Procédé selon la revendication 4, dans lequel le composé de potassium est le benzoate de potassium.
8. (S)-(+)-N,N-diméthyl-3-(1-naphtalényloxy)-3-(2-thiényl)propanamine, sel d'acide phosphorique.

30

35

40

45

50

55